Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

NOTE. Throughout the foregoing investigation it has been assumed that the signs of b_2, b_4, and b_6 in formulas (6), (7) and (8) will be +, — and + respectively, so that the signs of the constant a will be —, + and — respectively. This is true for any anjustment formula likely to be used in practice, but others can be constructed so that b_2, b_4 and b_6 will have signs contrary to the above. In such cases, the successive coefficients of the resultant formula tend to diverge from eachother as k increases, and in passing to the limit from (5) we cannot neglect differences of a higher order in comparison with those of a lower one. Take for example

$$u'_0 = \tfrac{1}{3}[3u_0+(u_1+u_{-1})+(u_2+u_{-2})],$$

for which $b_2 = -1$. The following table shows a few of the coefficients of the resultant formulas for 2, 4, 8 and 16 applications, as found by formula (7). (ANALYST, p. 67.) It is evident that, at the limit, some of the coefficients will become infinite and positive, others infinite and negative, and the excess of the sum of the positve over the sum of the negative ones will be unity. The series will be discontinuous, and there can be no limiting curve. Thus in (10), when b_2 is negative, h becomes imaginary; and in (13) and (19), if b_4 is +, then a becomes —, and A and B are imaginary.

	1	2	4	8	16
l_0	1.00	1.44	3.22	21.1	1297
l_1	.33	.44	.99	5.9	341
l_2	—.33	—.55	—1.73	—14.5	—1014
l_3		—.22	—1.04	—10.3	— 765
l_4		.11	.43	4.8	433
l_5			.35	6.7	717
l_6			— .07	— .4	— 4
&c.			&c.	&c.	&c.

NOTE ON THE CORRESPONDENCE OF MATERIAL FORMS WITH MATHEMATICAL RELATIONS.

BY THE EDITOR.

LIFE, sensation and consciousness are conditions found only in connection with definite material forms which constitute a part of the universe of matter; and the names used to denote the sensations experienced by the conscious individual are a representation of the external world. All knowledge is, therefore, but experience of impressions made by the world